

AD-A079 724

WISCONSIN UNIV-MADISON MATHEMATICS RESEARCH CENTER F/G 12/1  
ON THE MAXIMALITY OF THE SUM OF TWO MAXIMAL MONOTONE OPERATORS.(U)  
NOV 79 H ATTOUCH  
MRC-TSR-2018 DAAG29-75-C-0024  
NL

UNCLASSIFIED

1 OF 1  
AD-  
A079724




END  
DATE  
FILMED

2-80  
DDC









ADA 079724

9  
MRC TECHNICAL SUMMARY REPORT #2018

6  
ON THE MAXIMALITY OF THE SUM  
OF TWO MAXIMAL MONOTONE OPERATORS 3

10  
Hedy/Attouch

LEVEL

14  
Mathematics Research Center  
University of Wisconsin-Madison  
610 Walnut Street  
Madison, Wisconsin 53706

November 1979

DDC  
RECEIVED  
JAN 23 1980  
E

DDC FILE COPY

(Received August 15, 1979)

12/13  
15 DAAG29-75-C-~~44~~24

Approved for public release  
Distribution unlimited

Sponsored by

U. S. Army Research Office  
P. O. Box 12211  
Research Triangle Park  
North Carolina 27709

221 200

LB



UNIVERSITY OF WISCONSIN - MADISON  
MATHEMATICS RESEARCH CENTER

ON THE MAXIMALITY OF THE SUM OF TWO  
MAXIMAL MONOTONE OPERATORS

Hedy Attouch

Technical Summary Report #2018

November 1979

ABSTRACT

We prove that the topological condition  $\text{Int}(D(A) - D(B)) \neq \emptyset$ , implies that the sum of the two maximal monotone operators  $A$  and  $B$  is still maximal monotone.

AMS (MOS) Subject Classifications: 47H05, 49A40


Key Words: Maximal monotone operators, Duality methods in optimization,  
Convexity

Work Unit Number 1 - Applied Analysis

This document has been approved  
for public release and sale; its  
distribution is unlimited.

Sponsored by the United States Army under Contract No. DAAG29-75-C-0024.

- i -





## SIGNIFICANCE AND EXPLANATION

A wide variety of problems involving nonlinear partial differential equations, subject to boundary conditions, may be shown to have solutions by establishing that the associated differential operators satisfy a certain technical condition.

This condition, called maximal monotonicity, allows the use of a well developed abstract theory which includes results about existence, regularity, etc., of solutions of operator equations.

It is quite useful, therefore, to have easily verified conditions which imply that an operator is maximal monotone.

Frequently an operator may be regarded as the sum of simpler components. This paper gives a new sufficient condition which guarantees that the sum of two maximal monotone operators is again maximal monotone.

Accession For	
NTIS GMA&I	<input checked="checked" type="checkbox"/>
DDC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or special
A	

The responsibility for the wording and views expressed in this descriptive summary lies with MRC, and not with the author of this report.



ON THE MAXIMALITY OF THE SUM OF TWO  
MAXIMAL MONOTONE OPERATORS<sup>†</sup>

Hedy Attouch

Introduction

Let  $A$  and  $B$  be two maximal monotone operators in a real Hilbert space  $H$ . The classical theorem of R. T. Rockafellar [7], H. Brezis [2] tells us that if  $D(A) \cap \text{Int } D(B) \neq \emptyset$ , then the sum  $A + B$  is still maximal monotone. We show that the same conclusion holds under the weaker, symmetric assumption:  $\text{Int}(D(A) - D(B)) \supsetneq \emptyset$  (Theorem 1)<sup>\*</sup>. When one of the operators is a subdifferential,  $A = \partial\varphi$ , (resp. when the two operators are subdifferentials  $A = \partial\varphi$ ,  $B = \partial\psi$ ) this condition can be weakened to  $\text{Int}(D(\varphi) - D(B)) \supsetneq \emptyset$  (Theorem 2) (resp.  $\text{Int}(D(\varphi) - D(\psi)) \supsetneq \emptyset$  (Theorem 3)). These results are intimately related (Remark 2 and Remark 3) to:

a) H. Brezis and A. Haraux in [4] give sufficient conditions on two maximal monotone operators  $A$  and  $B$ , in order that  $R(A+B) \subseteq R(A) + R(B)$  (i.e. that the range of the sum and the sum of the ranges have same interior and closure).

b) The classical "Slater stability condition" (Continuity of a functional at one point of the domain of the other) can be weakened (cf. for example, J. P. Aubin [1]) to the following symmetric one: the difference of their domains is a neighbourhood of the origin.

I thank very much J. P. Aubin and H. Brezis for their advise and stimulating discussions.

---

<sup>†</sup> These notes have been written while visiting the Mathematics Research Center of the University of Wisconsin-Madison (June-August 1979) during the CNRS-NSF visiting program G.05.0252.

---

<sup>\*</sup> In fact (cf. Remark 2) one can weaken this assumption to  $\text{Int}(\text{Conv } D(A) - \text{Conv } D(B)) \supsetneq \emptyset$ .



Let  $H$  be a real Hilbert space with the norm  $|\cdot|$ , and the scalar product  $\langle \cdot, \cdot \rangle$ . Given  $A$  a maximal monotone operator in  $H$ , we shall denote by  $D(A)$  its domain, by  $R(A)$  its range, by  $A_\lambda = \frac{1}{\lambda}(I - (I + \lambda A)^{-1})$ , for  $\lambda > 0$ , its Yosida approximation and by  $A^0$  its minimal section.

### Theorem 1

Let  $A$  and  $B$  two maximal monotone operators in  $H$ .

Let us assume that  $0 \in \text{Int}(D(A) - D(B))$ . (\*)

Then  $A + B$  is a maximal monotone operator.

### Proof of Theorem 1

Given  $f$  in  $H$  and  $\lambda > 0$ , let us consider  $u_\lambda$  the solution of

$$(1_\lambda) \quad u_\lambda + Au_\lambda + B_\lambda u_\lambda \ni f$$

and let us prove that  $\sup_{\lambda > 0} |B_\lambda u_\lambda| < +\infty$ ; from the Brezis/Crandall/Pazy theorem

[3] it will follow that the  $u_\lambda$  converges, as  $\lambda$  goes to zero, to a solution  $u$  of

$$(1) \quad u + Au + Bu \ni f,$$

and this implies the maximality of  $A + B$ .

We first observe that the  $(u_\lambda)_{\lambda > 0}$ , are bounded in  $H$ : taking  $x \in D(A) \cap D(B)$ , and using the monotonicity of  $A + B_\lambda$ , one has

$$|u_\lambda - x| \leq |f - (x + A^0 x + B_\lambda x)| \quad \text{which implies} \quad |u_\lambda| \leq 2|x| + |f| + |A^0 x| + |B^0 x|.$$

By assumption, there exists a  $\rho > 0$  such that

$$\forall y \in H, |y| < \rho \quad y \in D(A) - D(B).$$

Given such a  $y$ , it suffices to show that  $\sup_{\lambda > 0} \langle B_\lambda u_\lambda, y \rangle \leq C(y) < +\infty$ ; the conclusion will then follow from the uniform boundedness theorem.

If  $y \in H$ ,  $y$  can be written as  $y = \beta - \alpha$  with  $\beta \in D(B)$ ,  $\alpha \in D(A)$ , and

---

(\*) That is to say, the algebraic difference of their domains is a neighbourhood of the origin.



$$\langle B_\lambda u_\lambda, y \rangle = \langle B_\lambda u_\lambda, \beta \rangle - \langle B_\lambda u_\lambda, \alpha \rangle .$$

a) From the monotonicity of  $B_\lambda$ ,

$$\langle B_\lambda u_\lambda - B_\lambda \beta, u_\lambda - \beta \rangle \geq 0, \text{ which implies}$$

$$\langle B_\lambda u_\lambda, \beta \rangle \leq \langle B_\lambda u_\lambda, u_\lambda \rangle - \langle B_\lambda \beta, u_\lambda - \beta \rangle ;$$

$$\text{so, } \langle B_\lambda u_\lambda, y \rangle \leq \langle B_\lambda u_\lambda, u_\lambda - \alpha \rangle + |B^0 \beta| \cdot |u_\lambda - \beta| .$$

b) Since  $u_\lambda$  satisfies  $(1_\lambda)$ ,  $B_\lambda u_\lambda = f - u_\lambda - v_\lambda$  with  $v_\lambda \in Au_\lambda$ , and the preceding inequality becomes:

$$\langle B_\lambda u_\lambda, y \rangle \leq \langle f - u_\lambda - v_\lambda, u_\lambda - \alpha \rangle + |B^0 \beta| \cdot |u_\lambda - \beta| .$$

c) From the monotonicity of  $A$ ,

$$\langle v_\lambda - A^0 \alpha, u_\lambda - \alpha \rangle \geq 0 \Rightarrow - \langle v_\lambda, u_\lambda - \alpha \rangle \leq |A^0 \alpha| \cdot |u_\lambda - \alpha| .$$

Finally,

$$\langle B_\lambda u_\lambda, y \rangle \leq |u_\lambda - \alpha| [|f - u_\lambda| + |A^0 \alpha|] + |u_\lambda - \beta| \cdot |B^0 \beta| \leq C(y)$$

since the  $(u_\lambda)$  are bounded for  $\lambda > 0$ .

Remark 1. The condition  $\text{Int}(D(A) - D(B)) \supset 0$ , is symmetric in  $A$  and  $B$  and is weaker than the classical condition of Rockafellar [7], H. Brezis [2],  $\text{Int } D(A) \cap D(B) \neq \emptyset$ , since  $\text{Int}(D(A) - D(B)) \supset \text{Int } D(A) - D(B)$ . Clearly this weaker condition can be satisfied even if  $D(A)$  and  $D(B)$  have an empty interior domain!

Let us see now how to improve this result when one of the operator is a sub-differential.

#### Theorem 2

Let  $B$  be a maximal monotone operator in  $H$  and  $\partial \varphi$  be the subdifferential of  $\varphi$ , a convex, lower semi continuous, proper function from  $H$  into  $]-\infty, +\infty]$ .

Let us assume that  $0 \in \text{Int}(D(\varphi) - D(B))$

Then  $\partial \varphi + B$  is a maximal monotone operator.



### Proof of Theorem 2

As in the proof of Theorem 1, we consider the solution  $u_\lambda$  of

$$(2_\lambda) \quad u_\lambda + \partial\varphi(u_\lambda) + B_\lambda u_\lambda \ni f.$$

We want to prove that  $\sup_\lambda |B_\lambda u_\lambda| < +\infty$ . The equation  $(2_\lambda)$  can be written

$$(2_\lambda)_{bis} \quad \forall v \in H \quad 0 \leq \varphi(u_\lambda) - \varphi(v) + \langle f - u_\lambda - B_\lambda u_\lambda, v - u_\lambda \rangle.$$

The  $(u_\lambda)_{\lambda>0}$  are bounded in  $H$ : Let us take  $v_0 \in D(\varphi) \cap D(B)$  and  $b, c \in \mathbb{R}^+$  such that  $\varphi(x) + b|x| + c \geq 0$ ; Since  $\langle B_\lambda u_\lambda - B_\lambda v_0, u_\lambda - v_0 \rangle \geq 0$ , from  $(2_\lambda)_{bis}$  we get:

$$-b|u_\lambda| - c - \varphi(v_0) + \langle f - u_\lambda, v_0 - u_\lambda \rangle + \langle B_\lambda v_0, u_\lambda - v_0 \rangle \leq 0;$$

so there exist  $p_1$  and  $p_2 \in \mathbb{R}$  such that:  $\forall \lambda > 0 \quad |u_\lambda|^2 + p_1|u_\lambda| + p_2 \leq 0$ , which implies that the  $(u_\lambda)$  are bounded for  $\lambda > 0$ .

By assumption, there exists a  $\rho > 0$  such that

$$\forall y \in H \quad |y| < \rho \quad y \in D(\varphi) - D(B).$$

Let us prove that for any such  $y$

$$\sup_\lambda \langle B_\lambda u_\lambda, y \rangle \leq C(y) < +\infty,$$

the conclusion will follow from the uniform boundedness theorem; so  $y = \beta - \alpha$ ,  $\beta \in D(B)$ ,  $\alpha \in D(\varphi)$  and

$$\langle B_\lambda u_\lambda, y \rangle = \langle B_\lambda u_\lambda, \beta \rangle - \langle B_\lambda u_\lambda, \alpha \rangle.$$

From the monotonicity of  $B_\lambda$

$$\langle B_\lambda u_\lambda, y \rangle \leq \langle B_\lambda u_\lambda, u_\lambda - \alpha \rangle + |B^0 \beta| \cdot |u_\lambda - \beta|.$$

Now take  $v = \alpha$  in  $(2_\lambda)_{bis}$ :

$$\langle B_\lambda u_\lambda, u_\lambda - \alpha \rangle \leq \varphi(\alpha) - \varphi(u_\lambda) + \langle f - u_\lambda, u_\lambda - \alpha \rangle.$$

Combining the two last inequalities, we get



$$\langle B_\lambda u_\lambda, y \rangle \leq \varphi(\alpha) + b|u_\lambda| + c + |f - u_\lambda| \cdot |u_\lambda - \alpha| + |B^0 \beta| \cdot |u_\lambda - \beta|$$

$\leq C(y)$  since the  $(u_\lambda)_{\lambda > 0}$  are bounded.

Finally, when the two operators are subdifferentials we obtain the following result: We denote by  $\varphi^*$  the conjugate function of  $\varphi$  and by  $\nabla$  the inf convolution (i.e. given two functions  $\varphi_1$  and  $\varphi_2$ , then  $\varphi_1 \nabla \varphi_2(x) = \inf_y \{ \varphi_1(x-y) + \varphi_2(y) \}$ ).

### Theorem 3

Let  $\varphi$  and  $\psi$  two convex, lower semi-continuous, proper functions from  $H$  into  $]-\infty, +\infty]$ .

Let us assume that  $0 \in \text{Int}(D(\varphi) - D(\psi))$ .

Then  $\varphi^* \nabla \psi^*$  is lower semi continuous, the inf-convolution is exact and  $\partial\varphi + \partial\psi$  is a maximal monotone operator.

### Proof of Theorem 3

a) Let  $c \in \mathbb{R}$  and  $f_n \xrightarrow{s-H} f$  such that  $\varphi^* \nabla \psi^*(f_n) \leq c$ ; let us prove that  $\varphi^* \nabla \psi^*(f) \leq c$ ; from the definition of the inf-convolution, there exist  $\varepsilon_n \rightarrow 0$  and  $u_n \in H$  such that:

$$(3) \quad \varphi^*(u_n) + \psi^*(f_n - u_n) \leq C + \varepsilon_n.$$

Let us prove that  $\sup_n |u_n| < +\infty$ ; then taking a weakly converging subsequence  $u_{n_k} \xrightarrow{w-H} \bar{u}$ , by the lower semi-continuity of  $\varphi^*$  and  $\psi^*$ , we shall get

$$\varphi^*(\bar{u}) + \psi^*(f - \bar{u}) \leq C \text{ and therefore } \varphi^* \nabla \psi^*(f) \leq C.$$

The same argument will tell us that

$$\forall f \in D(\varphi^* \nabla \psi^*) \exists u \in H \text{ such that } \varphi^* \nabla \psi^*(f) = \varphi^*(u) + \psi^*(f - u)$$

(i.e. the inf-convolution is exact).

In order to prove the boundedness of the  $(u_n)_{n \in \mathbb{N}}$  we apply the uniform boundedness theorem: let  $\rho > 0$  such that



$$\forall y \in H \quad |y| < \rho \Rightarrow y \in D(\varphi) - D(\psi)$$

and let us prove that for any such  $y \sup_n \langle u_n, y \rangle \leq C(y) < +\infty$ .

So  $y = \alpha - \beta$ , with  $\alpha \in D(\varphi)$ ,  $\beta \in D(\psi)$  and

$$\langle u_n, y \rangle = \langle u_n, \alpha \rangle + \langle f - u_n, \beta \rangle - \langle f, \beta \rangle$$

$$\leq \varphi^*(u_n) + \varphi(\alpha) + \psi^*(f - u_n) + \psi(\beta) - \langle f, \beta \rangle.$$

From (3)

$$\langle u_n, y \rangle \leq C + \varepsilon_n + \varphi(\alpha) + \psi(\beta) - \langle f, \beta \rangle$$

and the  $(u_n)_{n \in \mathbb{N}}$  are bounded.

b) Let us take  $(u, f) \in \partial(\varphi + \psi)$  and let us prove that  $(u, f) \in \partial\varphi + \partial\psi$ ; from the maximal monotony of  $\partial(\varphi + \psi)$  the result will follow:

$$(u, f) \in \partial(\varphi + \psi) \iff (\varphi + \psi)(u) + (\varphi + \psi)^*(f) - \langle f, u \rangle = 0.$$

From a),  $(\varphi + \psi)^* = (\varphi^* \nabla \psi^*)^{**} = \varphi^* \nabla \psi^*$ , since  $\varphi^* \nabla \psi^*$  is lower-semi-continuous.

Moreover, there exists  $\bar{u} \in H$  such that  $(\varphi^* \nabla \psi^*)(f) = \varphi^*(\bar{u}) + \psi^*(f - \bar{u})$ . So,  $\{\varphi(u) + \varphi^*(\bar{u}) - \langle u, \bar{u} \rangle\} + \{\psi(u) + \psi^*(f - \bar{u}) - \langle u, f - \bar{u} \rangle\} = 0$ . Since each of these quantities is positive, each is equal to zero which means that

$$\bar{u} \in \partial\varphi(u), \quad f - \bar{u} \in \partial\psi(u) \quad \text{and} \quad f = \bar{u} + (f - \bar{u}) \in (\partial\varphi + \partial\psi)(u).$$

## Remark 2

The Brezis-Haraux theorem [4] states:

Let  $A$  and  $B$  two maximal monotone operators satisfying:

$A+B$  is maximal monotone

$A$  and  $B$  satisfy the condition (\*)

then  $R(A+B) \subseteq R(A) + R(B)$ .

We recall that a monotone operator  $S$  satisfies (\*) if

$$\forall x \in D(S) \quad \forall y \in D(S) \quad \exists C(x, y) \quad \text{such that:} \quad \forall z \in D(A) \quad \langle Sz - Sx, z - y \rangle \geq C(x, y).$$



As noticed by Brezis, one can deduce the theorem 1 from this theorem through the following transformation:

Let us consider the equation (1)  $2u + Au + Bu \ni f$ , which we want to solve.

Let us write it  $u + Au + u + Bu \ni f$ ; since  $I + B$  is onto, let us define  $v = u + Bu$  as a new variable; the equation to be solved becomes

$$(4) \quad (I+B)^{-1}(v) - (I+A)^{-1}(f-v) = 0.$$

We apply the B/H theorem with  $Sv = (I+B)^{-1}(v)$ ,  $Tv = -(I+A)^{-1}(f-v)$ . Clearly, these two operators are maximal monotone continuous operators and their sum is maximal monotone. They satisfy the condition (\*):

$$\begin{aligned} \langle (I+B)^{-1}(z) - (I+B)^{-1}(x), z-y \rangle &= \langle u-v, z-y \rangle \text{ with } u = (I+B)^{-1}z, v = (I+B)^{-1}x \\ &= \langle u-v, u + Bu - y \rangle \\ &= \langle u-v, u-v + (v+Bv) + (Bu-Bv) - y \rangle \\ &\geq |u-v|^2 - |x+y| \cdot |u-v| \geq C(x,y). \end{aligned}$$

$$\begin{aligned} \text{Therefore, } R((I+B)^{-1} - (I+A)^{-1}(f-)) &\supset \text{Int}[\text{Conv } R(I+B)^{-1} + \text{Conv } R(-(I+A)^{-1}(f-))] \\ &\supset \text{Int}[\text{Conv } D(B) - \text{Conv } D(A)]. \end{aligned}$$

So if we assume that  $\text{Int}[\text{Conv } D(B) - \text{Conv } D(A)] \ni 0$  we can solve (4) and therefore (1).

### Remark 3

The main part of the proof of the Theorem 3 is the following:

If  $0 \in \text{Int}(D(\varphi) - D(\psi))$  then (5)  $(\varphi+\psi)^* = \varphi^* \vee \psi^*$ . This result improves the classical result of Moreau [6]. In other words, the condition  $0 \in \text{Int}(D(\varphi)-D(\psi))$ , which is in fact a Slater-stability condition, (weaker than the classical one), implies the equality (5) of the primal and dual problem, and the existence of a solution for the dual problem. More generally, in the variational situation studied by Ekeland-Temam, [5], one can prove the following statement:



Theorem (See for instance J. P. Aubin [1] Ch. 14).

Let  $V$  and  $Y$  two reflexive Banach spaces and

$A : V \rightarrow Y$  a continuous linear map

$\varphi : Y \rightarrow ]-\infty, +\infty]$  a convex, lsc, proper function

$\psi : V \rightarrow ]-\infty, +\infty]$  a convex, lsc, proper functions

such that  $\text{Int}\{A^D(\psi) - D(\varphi)\} \neq \emptyset$ .

Then  $\forall f \in V^* \quad \inf_{u \in V} \{\psi(u) + \varphi(Au) - (f, u)\} = - \min_{g \in Y^*} \{\psi^*(f - A^*g) + \varphi^*(g)\}.$

Taking  $A = \text{Id}^+$  with  $V = Y$  we get the previous result.



## REFERENCES

- [1] Aubin, J. P., Mathematical Methods of Game and Economic Theory. North-Holland.
- [2] Brezis, H., Operateurs maximaux monotones et semi-groupes de contractions dans les espaces de Hilbert. North-Holland 1971.
- [3] Brezis, H., M. Crandall and A. Pazy, Perturbations of nonlinear maximal monotone sets in Banach spaces. Comm. Pure Appl. Math. 23, 123-144 (1970).
- [4] Brezis, H. and A. Haraux, Image d'une Somme d'Operateurs Monotones et applications. Israel Journal of Math. Vol. 23, No. 2, 1976.
- [5] Ekeland, I. and R. Temam, Analyse Convexe et problèmes variationnels. Dunod. Gauthier-Villars 1974.
- [6] Moreau, J. J., Fonctionnelles convexes. Seminaire sur les équations aux Dérivées partielles. Collège de France 1966-67.
- [7] Rockafellar, R. T., Conjugate duality and optimization. Philadelphia 1976. SIAM Coll. Regional conference series in applied mathematics No. 16.

HA/jvs



REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER #2018	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle)  ON THE MAXIMALITY OF THE SUM OF TWO MAXIMAL MONOTONE OPERATORS		5. TYPE OF REPORT & PERIOD COVERED Summary Report - no specific reporting period
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s)  Hedy Attouch		8. CONTRACT OR GRANT NUMBER(s)  DAAG29-75-C-0024
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of 610 Walnut Street Wisconsin Madison, Wisconsin 53706		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Work Unit Number 1 - Applied Analysis
11. CONTROLLING OFFICE NAME AND ADDRESS U. S. Army Research Office P. O. Box 12211 Research Triangle Park, North Carolina 27709		12. REPORT DATE November 1979
		13. NUMBER OF PAGES 9
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report)  UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)  Maximal monotone operators, Duality methods in optimization, Convexity.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)  We prove that the topological condition $\text{Int } (D(A) - D(B)) \neq \emptyset$ , implies that the sum of the two maximal operators A and B is still maximal monotone.		